

Archives of Orofacial Sciences

print ISSN: 1823-8602

e-ISSN: 2231-7163

CASE REPORT

To appear in: Archives of Orofacial Sciences

Submitted: 17th February 2020

Accepted: 8th October 2020

Published online: 18th October 2020

This is a PDF file of an article that has undergone enhancements after acceptance and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 The Authors. Production and hosting by Penerbit USM, on behalf of Universiti Sains Malaysia.

Surgical Management for Symptomatic Huge Frontoethmoidal Osteoma: A Case Report

Che Mohd Hilmi Che Mat ^{a,b}, Rosli Mohd Noor ^b, Rosdi Ramli ^b, Irfan Mohamad ^{a*}

^a Department of Otorhinolaryngology-Head & Neck Surgery, School of Medical Sciences, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia.

^b Department of Otorhinolaryngology-Head & Neck Surgery, Hospital Raja Perempuan Zainab II, 15200 Kota Bharu, Kelantan, Malaysia.

* Corresponding author: irfankb@usm.my

How to cite this article: Che Mat CMH, Mohd Noor R, Ramli R, Mohamad I (2020). Surgical Management for Symptomatic Huge Frontoethmoidal Osteoma: A Case Report. *Arch Orofac Sci*.

Submitted: 17/02/2020. Accepted: 08/10/2020. Published online: 18/10/2020.

Abstract:

Osteomas of paranasal sinuses are slow-growing benign tumour which is often asymptomatic. The prevalence is highest amongst fourth and sixth decades. Paranasal sinus osteomas are most commonly found in the frontal sinus. Osteomas are usually asymptomatic and found incidentally on radiographic examination. The symptoms depend on its size and location. Common symptoms include frontal pain, nasal discharge, posterior nasal drip, bump over the inner aspect of the lower eyelid and epiphora. We described a case of 34-year-old woman with an enlarging right medial canthal swelling for ten years, with increased intercanthal distance followed by persistent right nasal blockage. Because of worsening symptoms, she had sought for treatment in which endoscopic approach excision was performed.

Keyword: Benign Bony Tumour; Endoscopy, Osteoma; Paranasal Sinus; Surgery.

Introduction:

Osteoma is a benign osteogenic tumour. It arises from the compact or cancellous bone (Bulut *et al.*, 2010). It is common in the paranasal sinus, involving ethmoid and frontal sinus in 95% of cases (Cheng *et al.*, 2013). Osteomas are often seen in males (Kaplan *et al.*, 1994). It grows gradually and can produce facial asymmetry and other space occupying symptoms in the nose and paranasal sinuses. Osteomas are usually asymptomatic and found incidentally on radiographic examination. The incidence of frontal osteoma was only 0.4% from the reviewed asymptomatic 3,500 skull radiographs (Childrey, 1939). In some cases, osteoma may produce symptoms related to the surrounding structures.

Case summary:

A 34-year-old woman complained of swelling at right medial canthus for more than 10 years. It was associated with right epiphora, increased in intercanthal distance and persistent right nasal blockage. There was no epistaxis, nasal discharge, facial discomfort, or frontal pain. Examination of the face revealed a swelling over the right medial canthus, measuring 2 cm x 2 cm. The mass was firm, non-mobile with no sign of inflammation.

Nasoendoscopy showed multilobulated bony-hard mass pushing the inferior turbinate and septum laterally. The mass was insensitive to touch. Biopsy was not possible due to its hard consistency. A computed tomography (CT) scan of paranasal sinus revealed a well-circumscribed intra-sinus calcified mass occupying the right frontal and ethmoid sinus as well as right inferior meatus (Figure 1). Its measurement was 2.6 cm x 2.3 cm x 5.0 cm.

There was also an expansion of right ethmoid sinus causing the remodelling of the medial wall of the right orbit. The right nasolacrimal duct was compressed. Right fronto-ethmoidal recess was obliterated. Right osteo-meatal complex was preserved. Mucosal thickening of bilateral maxillary sinuses was noted. Left frontal and sphenoid sinuses were normal. Nasal septum was deviated to the left (Figure 2). Visualized brain was normal with no focal enhancing parenchymal lesion. Radiological impression was right-sided intranasal calcified mass. The differential diagnosis included chronic mycetoma, less likely a calcified foreign body.

The patient underwent endoscopic excision of right intranasal osteoma. Intraoperatively there was a bony mass occupying the whole right nasal cavity (Figure 3). The intact septum was pushed to the left side, with left middle turbinate and inferior turbinate lateralized. The mass was removed completely at the lower part and partly from the superior part. As tumour was very huge occupying the whole space in the nasal cavity, it was only managed to be removed in pieces using drill, chisel and mallet (Figure 4). Posterior part of inferior turbinate, middle part of middle turbinate and posterior choana were clearly visualized after removal of the mass. Merocele was inserted and oral antibiotic was started post operatively. Patient was discharged on the next day.

At one week follow-up after surgery, the patient claimed there was no epiphora, epistaxis, or nasal blockage. Nasoendoscopy showed minimal crusting. Histopathological examination confirmed an osteoma. Microscopically showed fibroadipose and fibrocollagenous tissue with dense lamellar bone having haversian canals and medullary component coexisting of fibrous tissue. The stroma was congested with moderate infiltrations by lymphoplasmacytic cells. No evidence of atypia or malignancy seen.

Discussion:

The most common benign tumours of the paranasal sinuses are osteoma. The occurrence is around 0.5% with men being slightly more frequently affected (Mesoella *et al.*, 2005). The aetiology includes hereditary, embryonic malformation, inflammation, trauma, calcification of polyposis and calcium metabolism disorder. They are commonly found in the frontal sinus. The tumour grows slowly over time which may be symptomatic in 10% of cases (Furlaneto *et al.*, 2004).

The symptoms are dependent on its location and size. Drainage and ventilation of paranasal sinuses can be affected by ingrowing osteoma which distorted the anatomy of paranasal sinuses. In the frontal sinus, the patient might complaint of frontal headache, frontal pain, sensation of pressure, protrusion of anterior wall frontal sinus which can lead to cosmetically problematic. If osteomas extend beyond the sinus margin, it will cause complications like intracranial mucocoeles, meningitis and abscesses (Al-Sebeih and Desrosiers, 1998). In case of sphenoid sinus osteomas, surgical removal should be done immediately because of the probable danger of the rapid compression on the visual pathways that leads to blindness (Mansour *et al.*, 1999).

Osteomas in fronto-ethmoidal complex may cause headache (49%), proptosis (40%), diplopia (18%) and hypoesthesia (13%). In ethmoidal osteoma, the ocular symptoms were much more common (66%) while in frontal osteoma, the headache was much more common (61%) (Cheng *et al.*, 2013). Giant frontal osteoma predominantly causes intracranial complication (54%) while giant ethmoidal osteoma causes more intraorbital complication (69%) (Cheng *et al.*, 2013). In our case, the patient did not complain of any intracranial or ocular symptoms. The main symptoms in our patient were right epiphora and right nasal blockage.

CT scan is the excellent imaging modality for the diagnosis of osteomas. It specifies the anatomical location and the tumour extension. In case of incomplete resection, the recurrence rate was 10% and no malignant transformation has been reported (Grayeli *et al.*, 1998)

In asymptomatic osteoma, the surgical intervention is not indicated (Castelnuovo *et al.*, 2008). There is no role for medical treatment in osteoma. In our case, surgical removal was indicated as she had the troublesome right epiphora and persistent right nasal blockage. The approach was either by external nasal, endoscopic or combination of both. We partly removed the osteoma by endoscopic approach by using telescope (0°, 30° and 70° angles of view) with few types of surgical instruments. We had difficulty to remove all the tumours as it was huge and extending to the frontal sinus. We also anticipated the problem during the endoscopic removal as there was no adequate space for surgical instrument to enter the intranasal cavity. However, we managed to remove the lower part of osteoma at the level of middle turbinate to relieve the symptoms (nasal blockage and epiphora) and to avoid any external scar in this young female patient. She remained symptom-free during the one-month clinic review. Endoscopy assessment showed no evidence of regrowth of the mass. The follow-up was done for six months.

Frontal sinus osteoma is staged according to system proposed by Chiu *et al.* (2005) (Table 1). In stage 1 and 2, the operation can be done by endoscopic surgery while in stage 3 and 4 by external approach. The external approach gives the surgeon a wider vision, but the disadvantages are post operation oedema, pain and incision scars. Frontal ethmoidectomy (Lynch procedure) and osteoplastic flap can be done in external approach (Yiotakis *et al.*, 2008). Our patient was categorized as stage 4, as the tumour had already filled up the whole frontal sinus.

Conclusion:

Even though osteoma is a slow growing benign tumour, symptomatic paranasal sinus osteoma needs surgical excision. The extent of removal is a multi-factor dependant decision. Avoiding facial scar for cosmetic reason in a young woman is important as long as the symptoms are eradicated. CT scan is the best imaging modality to look for the site and extension of tumour. Endoscopic approach offers safe removal of tumour with better cosmetic outcomes. The external approach might be considered in the future if the tumour re-expands and symptoms recur.

Accepted Article

References:

- Al-Sebeih K, Desrosiers M (1998). Bifrontal endoscopic resection of frontal sinus osteoma. *Laryngoscope*, **108**(2): 295-298. <https://doi.org/10.1097/00005537-199802000-00026>
- Bulut E, Acikgoz A, Ozan B, Gunhan O (2010). Large peripheral osteoma of the mandible: A case report. *Int J Dent*, **2010**: 834761. <https://doi.org/10.1155/2010/834761>
- Castelnuovo P, Valentini V, Giovannetti F, Bignami M, Cassoni A, Iannetti G (2008). Osteomas of the maxillofacial district: Endoscopic surgery versus open surgery. *J Craniofac Surg*, **19**(6): 1446-1452. <https://doi.org/10.1097/SCS.0b013e31818b417d>
- Cheng KJ, Wang SQ, Lin L (2013). Giant osteomas of the ethmoid and frontal sinuses: Clinical characteristics and review of the literature. *Oncol Lett*, **5**(5): 1724-1730. <https://doi.org/10.3892/ol.2013.1239>
- Childrey JH (1939). Osteoma of the sinuses, the frontal and the sphenoid bone: Report of fifteen cases. *Arch Otolaryngol*, **30**(1): 63-72. <https://doi.org/10.1001/archotol.1939.00650060071006>
- Chiu AG, Schipor I, Cohen NA, Kennedy DW, Palmer JN (2005). Surgical decisions in the management of frontal sinus osteomas. *Am J Rhinol*, **19**(2): 191-197. <https://doi.org/10.1177/194589240501900213>
- Furlaneto E, Rocha JRM, Heitz C (2004). Osteoma of the zygomatic arch: Report of a case. *Int J Oral Maxillofac Surg*, **33**(3): 310-311. <https://doi.org/10.1006/ijom.2002.0468>
- Grayeli AB, Redondo A, Sterkers O (1998). Anterior skull base osteoid osteoma: Case report. *Br J Neurosurg*, **12**(2): 173-175. <https://doi.org/10.1080/02688699845366>
- Kaplan I, Calderon S, Buchner A (1994). Peripheral osteoma of the mandible: A study of 10 new cases and analysis of the literature. *J Oral Maxillofac Surg*, **52**(5): 467-470. [https://doi.org/10.1016/0278-2391\(94\)90342-5](https://doi.org/10.1016/0278-2391(94)90342-5)
- Mansour AM, Salti H, Uwaydat S, Dakroub R, Bashshour Z (1999). Ethmoid sinus osteoma presenting as epiphora and orbital cellulitis. *Surv Ophthalmol*, **43**(5): 413-426. [https://doi.org/10.1016/s0039-6257\(99\)00004-1](https://doi.org/10.1016/s0039-6257(99)00004-1)
- Mesolella M, Galli V, Testa D (2005). Inferior turbinate osteoma: A rare cause of nasal obstruction. *Otolaryngol Head Neck Surg*, **133**(6): 989-991. <https://doi.org/10.1016/j.otohns.2005.03.045>
- Yiotakis I, Eleftheriadou A, Giotakis E, Manolopoulos L, Ferekidou E, Kandiloros D (2008). Resection of giant ethmoid osteoma with orbital and skull base extension followed by duraplasty. *World J Surg Oncol*, **6**: 110. <https://doi.org/10.1186/1477-7819-6-110>




Figure 1: Coronal CT showed a well-circumscribed calcified mass occupying the right frontal sinus, ethmoid sinus and right nasal cavity. Red arrow: maxillary sinus. Green arrow: orbital rim. Blue arrow: vomer. Orange arrow: crista galli. Yellow arrow: base of skull.



Figure 2: Axial CT paranasal sinus revealed nasal septal deviation to the left.

Accepted Article




Figure 3: Intraoperative nasoendoscopic view showed multilobulated whitish bony mass in right nasal cavity.



Figure 4: Osteoma removed in pieces with the largest was 4.5 cm.

Table 1: The frontal sinus osteoma staging system by Chiu *et al.* (2005)

Stage	Criteria
Stage 1	Source of the osteoma posterior and inferior in the frontal recess; localization of the osteoma medial to a virtual sagittal plane passing over the lamina papyracea; anteroposterior (AP) diameter of the tumour is fewer than 75% of the AP diameter of the frontal sinus
Stage 2	As stage 1; AP diameter of the tumour larger than 75% of the AP diameter of the frontal recess
Stage 3	Source of the osteoma anterior and/or superior in the frontal sinus and/or osteoma spreading lateral to a virtual sagittal plane passing through the lamina papyracea
Stage 4	The tumour filling the whole frontal sinus

Accepted Article